

## Aberystwyth University

### *High C3 photosynthetic capacity and high intrinsic water use efficiency underlies the high productivity of the bioenergy grass Arundo donax*

Webster, Richard John; Driever, Steven; Kromdijk, Wanne; McGrath, Justin; Leakey, Andrew; Siebke, Katharina; Demetriades-Shah, Tanvir; Bonnage, Steven; Lawson, Tracy; Long, Stephen

*Published in:*  
Scientific Reports

*DOI:*  
[10.1038/srep20694](https://doi.org/10.1038/srep20694)

*Publication date:*  
2016

*Citation for published version (APA):*

Webster, R. J., Driever, S., Kromdijk, W., McGrath, J., Leakey, A., Siebke, K., Demetriades-Shah, T., Bonnage, S., Lawson, T., & Long, S. (2016). High C3 photosynthetic capacity and high intrinsic water use efficiency underlies the high productivity of the bioenergy grass *Arundo donax*. *Scientific Reports*, 6, [20694].  
<https://doi.org/10.1038/srep20694>

#### **Document License** CC BY

#### **General rights**

Copyright and moral rights for the publications made accessible in the Aberystwyth Research Portal (the Institutional Repository) are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Aberystwyth Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Aberystwyth Research Portal

#### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

tel: +44 1970 62 2400  
email: [is@aber.ac.uk](mailto:is@aber.ac.uk)

# SCIENTIFIC REPORTS

OPEN

## High C3 photosynthetic capacity and high intrinsic water use efficiency underlies the high productivity of the bioenergy grass *Arundo donax*

Received: 08 October 2015  
Accepted: 18 December 2015  
Published: 10 February 2016

Richard J. Webster<sup>1</sup>, Steven M. Driever<sup>2</sup>, Johannes Kromdijk<sup>4</sup>, Justin McGrath<sup>4</sup>, Andrew D. B. Leakey<sup>3,4</sup>, Katharina Siebke<sup>5</sup>, Tanvir Demetriades-Shah<sup>6</sup>, Steve Bonnage<sup>7</sup>, Tony Peloe<sup>8</sup>, Tracy Lawson<sup>9</sup> & Stephen P. Long<sup>3,4</sup>

*Arundo donax* has attracted interest as a potential bioenergy crop due to a high apparent productivity. It uses C3 photosynthesis yet appears competitive with C4 grass biomass feedstock's and grows in warm conditions where C4 species might be expected to be that productive. Despite this there has been no systematic study of leaf photosynthetic properties. This study determines photosynthetic and photorespiratory parameters for leaves in a natural stand of *A. donax* growing in southern Portugal. We hypothesise that *A. donax* has a high photosynthetic potential in high and low light, stomatal limitation to be small and intrinsic water use efficiency unusually low. High photosynthetic rates in *A. donax* resulted from a high capacity for both maximum Rubisco ( $V_{c,max}$   $117 \mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ ) and ribulose-1:5-bisphosphate limited carboxylation rate ( $J_{max}$   $213 \mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ ) under light-saturated conditions. Maximum quantum yield for light-limited  $\text{CO}_2$  assimilation was also high relative to other C3 species. Photorespiratory losses were similar to other C3 species under the conditions of measurement (25%), while stomatal limitation was high (0.25) resulting in a high intrinsic water use efficiency. Overall the photosynthetic capacity of *A. donax* is high compared to other C3 species, and comparable to C4 bioenergy grasses.

Giant reed (*Arundo donax* L.) has attracted interest as a potential bioenergy crop, due to a high apparent productivity and suitability as an accepted feedstock for cellulosic fuel production<sup>1</sup>. It is found throughout the Mediterranean climate zones of the world and has been cultivated in Asia, southern Europe and North Africa for over a thousand years. It is a rhizomatous perennial herbaceous grass that produces stems of ca. 2–3 cm diameter that may grow to heights of 3–6 m. It is found in moist grasslands and wetlands although it is able to thrive across a wide range of soil types, while also being tolerant of mild drought and salinity, and is tolerant of water-logged soils<sup>2–4</sup>. In climates with warm winters it is evergreen but dies back in climates with freezing winter temperatures. Although there are related species within the genus, molecular analysis suggests that *A. donax* is monophyletic in origin and its spread has not involved hybridisation with the related species. These analyses also suggest that it originated in west Asia and then spread to the Mediterranean region<sup>5</sup>. The high productivity of *A. donax* is achieved in warm growing conditions despite the fact that it uses C3 photosynthesis, which might be expected to be a competitive disadvantage compared to C4 species under such growing conditions.

<sup>1</sup>Institute of Biological Environmental and Rural Sciences, Aberystwyth University, Aberystwyth, U.K. <sup>2</sup>Centre for Crop Systems Analysis, Wageningen University, Wageningen, the Netherlands. <sup>3</sup>Department of Plant Biology, University of Illinois at Urbana-Champaign, Urbana, Illinois, USA. <sup>4</sup>Carl R. Woese Institute for Genomic Biology, University of Illinois at Urbana-Champaign, Urbana, Illinois, USA. <sup>5</sup>Heinz Walz GmbH, Effeltrich, Germany. <sup>6</sup>LI-COR Biosciences UK Ltd., Cambridge, UK. <sup>7</sup>ADC BioScientific Ltd., Hoddesdon, UK. <sup>8</sup>Delta-T Devices Ltd., Cambridge, UK. <sup>9</sup>School of Biological Sciences, University of Essex, Wivenhoe Park, Colchester, U.K. Correspondence and requests for materials should be addressed to R.J.W. (email: rcw@aber.ac.uk)

Side-by-side trials in central Italy over 11 years showed an average yield of above-ground dry biomass of 28.7 t ha<sup>-1</sup> yr<sup>-1</sup> for the highly productive C4 perennial grass *Miscanthus x giganteus* Deu et Greef, but 37.7 t ha<sup>-1</sup> yr<sup>-1</sup> for *A. donax*<sup>6</sup>. Although this appears a relatively low yield for *M. x giganteus* under irrigated conditions in the Mediterranean<sup>7</sup>, the reported yield for *A. donax* is high by any measure. *A. donax* in central and southern Italy showed average dry matter yield over 3 years of up to 74 t ha<sup>-1</sup> yr<sup>-1</sup>, which exceeds the highest yields reported for *M. x giganteus*<sup>8</sup>. Similarly, in a multi-year study in Alabama *A. donax* achieved an average yield of 35.5 t ha<sup>-1</sup> yr<sup>-1</sup> compared to 23.5 t ha<sup>-1</sup> yr<sup>-1</sup> for switchgrass (*Panicum virgatum* L.)<sup>9</sup>. In a comparison with C4 sweet sorghum (*Sorghum bicolor* L.) in northern Italy, light penetration into an *A. donax* canopy studied over a three year period was found to be 60% higher and the efficiency of conversion of intercepted radiation into biomass was ca. 60% higher. This might be explained by its more effective distribution of light through the canopy, if it is coupled with a similar efficiency of leaf photosynthesis<sup>10,11</sup>. However, yields may be considerably lower on marginal land. On a sandy loam with limited nutrient availability *A. donax* required three years to accumulate 20 t ha<sup>-1</sup>, and build a rhizome mass of 16 t ha<sup>-1</sup><sup>4</sup>, by contrast to *M. x giganteus* which has proved significantly more productive on marginal sites<sup>12</sup>. While *M. x giganteus* was found particularly vulnerable to damage by water-logging of soils during autumn and winter, *A. donax* appeared to thrive in these situations. This tolerance of water-logging also allows *A. donax* to spread along watercourses from rhizome fragments where *M. x giganteus* cannot<sup>13</sup>. Its ability to survive anaerobic soil may explain its ability to tolerate high levels of reduced ions of heavy metals<sup>14</sup>. *A. donax* is clearly a resilient, but also very productive species. However, there is little fundamental knowledge available on how it is able to realise this productivity. Understanding this demands knowledge of leaf photosynthesis as well as canopy microclimate.

Leaf photosynthetic rates of CO<sub>2</sub> uptake (*A*) measured in *A. donax* growing on an estuary in S. Africa were between 20 and 37 μmol m<sup>-2</sup> s<sup>-1</sup><sup>15</sup>. By combining modulated chlorophyll fluorescence and gas exchange, photorespiration was estimated to decrease *A* by 30%. Although water potential declined to -2.1 MPa around noon on sunny days, stomatal conductance (*g<sub>s</sub>*) and *A* changed little. Such low leaf water potential would normally be expected to at least cause partial stomatal closure in most species<sup>15</sup>. The rates of stomatal conductance reported are at the upper end of those found in other C3 species but not much greater than those of the productive C4 bio-energy grasses *M. x giganteus* and *P. virgatum*, as well as productive modern cultivars of maize (*Zea mays* L.)<sup>16,17</sup>.

In summary, *A. donax* achieves and exceeds the productivities normally associated with C4 perennial grasses in warm climates. For a plant which forms a high leaf area index (LAI) monocultures in nature and as a crop, it would be expected to have a high photosynthetic capacity at both high and low light intensity under warm conditions. The high photosynthetic capacity is needed in the context of a dense canopy where shaded leaves need to achieve high efficiency, as well as those in full sun, and where all leaves need to be efficient under the low light conditions of dawn, dusk and cloudy days. Yet it is a C3 species, growing in warm conditions where only a C4 species might be expected to be photosynthetically efficient<sup>18</sup>. The limited published data suggests that *A. donax* may indeed have high light-saturated leaf photosynthetic rates, which may also be supported by high leaf conductances. High leaf conductances will allow a higher CO<sub>2</sub> concentration at ribulose-1:5-bisphosphate carboxylase/oxygenase (Rubisco), so increasing the rate of carboxylation and decreasing photorespiration, but at the expense of water loss and water use efficiency<sup>19</sup>.

To better define the photosynthetic capacity of *A. donax* and understand how this may explain its high productivity this study aimed to define key *in vivo* properties. These were: *J<sub>max</sub>*, the maximum whole chain electron transport rate supporting ribulose-1:5-bisphosphate (RuBP) regeneration, *V<sub>c,max</sub>*, the maximum rate of carboxylation that can be supported by Rubisco, the light-saturated rate of CO<sub>2</sub> assimilation (*A<sub>sat</sub>*) and the maximum quantum yield of CO<sub>2</sub> assimilation (*Φ<sub>CO2,max</sub>*) defined by the initial slope of the response of *A* to absorbed photon flux (*αI*). Concurrent measurement of water vapour flux and modulated chlorophyll fluorescence were used to determine *g<sub>s</sub>*, stomatal limitation to CO<sub>2</sub> uptake (*l*), intercellular CO<sub>2</sub> concentration (*c<sub>i</sub>*), and the whole chain electron transport rate (*J*). These parameters were determined for leaves in a natural stand of *A. donax* growing in a dry valley in S. Portugal. These are used to test the hypotheses that *A. donax* has a high photosynthetic potential in both high and low light, that stomatal limitation may be unusually small and thus intrinsic leaf water use efficiency poor. The *in vivo* measures: *J<sub>max</sub>*, *V<sub>c,max</sub>*, *A<sub>sat</sub>* and *Φ<sub>CO2,max</sub>*, are key to parameterization of the steady-state biochemical model of C3 photosynthesis<sup>20</sup> which underlies most models of crop and ecosystem productivity<sup>21</sup>. This study therefore also serves as a key resource for parameterizing production models of this emerging crop.

## Materials and Methods

**Field location.** The study was undertaken on a naturally occurring *A. donax* stand located at Quinta de São Pedro - Centro de estudos (Armadas, 2815–786 Sobreda, Portugal 38°38'40.6"N 9°11'34.5"W, altitude 85 m). A mixed 5 acre site with a disturbed dry valley community of annuals, patches of native macchia and scrub, pine woodland, introduced Acacia woodland and a few planted ornamentals on a Cambisol soils<sup>22</sup>.

**Meteorological assessment.** Relative humidity, air temperature and irradiance were measured over the sampling period using a Weather Station (WS-GP2 with GP2 Data Logger, Delta-T Devices Ltd, Cambridge, UK) (Fig. S1). Relative Humidity and air temperature were used to calculate vapour pressure deficit (VPD), as a virtual channel within the GP2. Soil temperature and soil water content were measured within the stand over the sampling period using soil moisture sensors (SM300 with GP2 Data Logger, Delta-T Devices Ltd, Cambridge, UK).

**Stomatal Conductance and *In situ* Chlorophyll Fluorescence.** Measurements of stomatal conductance were made with a diffusion porometer (AP4, Delta-T Devices Ltd, Cambridge, UK). All measurements of conductance and photosynthesis were made on the most recently expanded leaf, as judged by ligule emergences. Readings were taken on several leaves, locating the porometer at three points along each leaf: 1) the base just above the ligule, 2) the midpoint between ligule to tip; and 3) the area just below the leaf tip. Both the

adaxial and abaxial surfaces were measured at these points at midday and mid-afternoon following the procedures of Monteith<sup>23</sup>, and Potter<sup>24</sup>. Modulated chlorophyll fluorescence was monitored to estimate the operating efficiency of PSII ( $F_q'/F_m'$ )<sup>25,26</sup> on fully emerged leaves at midday and late afternoon using a fluorometer (MONITORING-PAM, Heinz Walz GmbH, Effeltrich, Germany) (Fig. S1).

**Photosynthetic intercellular- $\text{CO}_2$  response curves.** Leaves were sampled pre-dawn from the *A. donax* stand. Leaves were excised just above the ligule, and immediately re-cut under water and then kept in cool low-light conditions ( $20^\circ\text{C}$  and  $20\ \mu\text{mol m}^{-2}\text{s}^{-1}$ ) until use. The response of net leaf  $\text{CO}_2$  uptake rate ( $A$ ) to external  $\text{CO}_2$  concentration ( $c_a$ ) was assessed on five different leaves. Leaves were placed in the leaf cuvette of a portable open gas exchange system (LCPro-SD with  $6.25\text{ cm}^2$  cuvette; ADC BioScientific Ltd. Herts, England; or LI6400XT with  $6\text{ cm}^2$  cuvette; LI-COR Biosciences, Lincoln, Nebraska, USA). Air temperature was controlled at  $25^\circ\text{C}$ , leaf temperature  $27^\circ\text{C}$  and VPD at  $0.96\text{ kPa}$ . To fully induce photosynthesis before measurements commenced, leaves were first allowed to equilibrate at a photon flux ( $I$ ) of  $500\ \mu\text{mol m}^{-2}\text{s}^{-1}$  and reference  $c_a$  of  $400\ \mu\text{mol mol}^{-1}$  until  $A$  had reached a stable value, the  $I$  was then increase to  $1500\ \mu\text{mol m}^{-2}\text{s}^{-1}$ . When  $A$  had reached a stable value at  $I = 1500\ \mu\text{mol m}^{-2}\text{s}^{-1}$   $c_a$  was then changed to the following levels in sequence, 300, 250, 200, 150, 100, 50, 400, 600, 900, 1200 and  $1500\ \mu\text{mol mol}^{-1}$ . The leaf remained at each level until a stable  $A$  could be determined. Potential Rubisco carboxylation ( $V_{c,\text{max}}$ ) and electron transport through photosystem II ( $J_{\text{max}}$ ) were determined from the responses of  $A_{\text{sat}}$  to  $c_i$ . For each leaf values of  $A$  located above the transition between Rubisco-limited and RuBP- or electron transport-limited photosynthesis in the  $A/c_i$  response were used to solve for  $J_{\text{max}}$  using the equations representing RuBP regeneration-limited  $A$ <sup>20</sup>. Values were temperature adjusted to  $25^\circ\text{C}$  using the temperature response equations of Bernacchi *et al.* (2001)<sup>27</sup> for the Rubisco-limited, and Bernacchi *et al.* (2003)<sup>28</sup> for the RuBP regeneration limited portions of the  $A$  vs  $c_i$  curves. Stomatal limitation ( $I$ ) at the current ambient  $c_a$  of  $400\ \mu\text{mol mol}^{-1}$  was derived using the method outlined in Long and Bernacchi (2003)<sup>29</sup>.

**Photosynthetic light response curves.** Sampling and measurements of  $A$  vs.  $I$  were performed in parallel with  $A$  vs.  $c_i$  curves. Leaves were placed in the leaf cuvette of an integrated open gas exchange system and modulated chlorophyll fluorometer (GFS-3000FL, LED-Array/PAM-Fluorometer 3055-FL and 3080-O2 Oxygen sensor. Heinz Walz GmbH, Effeltrich, Germany). Leaves were adapted to an incident photon flux of  $1500\ \mu\text{mol m}^{-2}\text{s}^{-1}$  prior to measurements with the  $[\text{CO}_2]$  at  $400\ \mu\text{mol mol}^{-1}$  and leaf temperature controlled at  $25^\circ\text{C}$ . The VPD of the air entering the gas exchange system was maintained between  $0.8$  and  $1.1\text{ kPa}$ . Photosynthetic light response curves were then obtained at ambient (21%) and low (2%) oxygen concentrations. Photon flux was varied in a step-wise manner, either starting from the photon flux at which photosynthesis had been induced, after a steady-state  $A$  was obtained or alternating between different photon fluxes, in each case waiting for a steady-state to be obtained. Photon flux levels were 2000, 1500, 1000, 700, 500, 350, 250, 150, 75,  $45\ \mu\text{mol m}^{-2}\text{s}^{-1}$ . At each light level, once a new steady state was reached gas exchange rates were recorded. Simultaneously,  $F_q'/F_m'$  was determined via modulated chlorophyll fluorescence and application of a saturating flash as previously described<sup>25,26,30</sup>.

The light response of  $\text{CO}_2$  assimilation rate is described by a four-parameter non-rectangular hyperbola, according to Marshall and Biscoe<sup>31</sup>:

$$A = \frac{\Phi_{\text{max}} I + A_{\text{sat}} - \sqrt{[(\Phi_{\text{max}} I + A_{\text{sat}})^2 - 4\theta \Phi_{\text{max}} I A_{\text{sat}}]}}{2\theta} - R_d \quad (1)$$

where  $A$  is the  $\text{CO}_2$  assimilation rate ( $\mu\text{mol m}^{-2}\text{s}^{-1}$ ),  $\Phi_{\text{max}}$  is the apparent maximum quantum yield ( $\text{mol CO}_2$  ( $\text{mol photons}^{-1}$ )),  $I$  is the incident photon flux ( $\mu\text{mol m}^{-2}\text{s}^{-1}$ ),  $A_{\text{sat}}$  is the  $\text{CO}_2$  assimilation rate at saturating photon flux ( $\mu\text{mol m}^{-2}\text{s}^{-1}$ ),  $\theta$  is the curve convexity (dimensionless) and  $R_d$  is mitochondrial respiration in the light ( $\mu\text{mol m}^{-2}\text{s}^{-1}$ ). The light compensation point (LCP) was determined as the incident photon flux ( $I$ ) where  $A = 0$  as predicted from the fitted curve.

The quantum yield of  $\text{CO}_2$  assimilation,  $\Phi_{\text{CO}_2}$ , was determined according to Genty<sup>30</sup>:

$$\Phi_{\text{CO}_2} = \frac{A + R_d}{\alpha I} \quad (2)$$

where  $A$  is corrected for respiratory loss ( $R_d$ ) and divided by the absorbed photon flux ( $\alpha I$ ), with  $\alpha$  as the leaf absorbance. The method used to determine absorbance under the chamber lighting conditions is described later.

The parameters  $\Phi_{\text{CO}_2}$  and  $F_q'/F_m'$  have been shown to be linearly related under conditions where photorespiration is suppressed<sup>30,32,33</sup>. Under ambient (21%) and low (2%) oxygen atmosphere, the relationship can be used to assess the possible operation of electron acceptors other than  $\text{CO}_2$ , e.g. photorespiration or active oxygen production, which are manifest as an increased  $F_q'/F_m'$  relative to its corresponding  $\Phi_{\text{CO}_2}$ <sup>34</sup>.

The relation of the quantum yield of  $\text{CO}_2$  assimilation ( $\Phi_{\text{CO}_2}$ ) after Valentini<sup>35</sup> to the operating efficiency of PSII ( $F_q'/F_m'$ ) under non-photorespiratory conditions can be described as a linear relationship<sup>30</sup>, where  $b$  is the  $y$  axis intercept and  $k$  is the slope of the relationship between  $F_q'/F_m'$  and  $\Phi_{\text{CO}_2}$ :

$$\frac{F'}{F'_m} = k * \Phi_{\text{CO}_2} + b \quad (3)$$

Assuming 4 electrons are necessary for the fixation of a  $\text{CO}_2$  molecule, this can be expressed as:

$$\frac{F'q}{F'm} = \frac{1}{4}k * \Phi_{e-} + b \quad (4)$$

where  $\Phi_{e-}$  is the quantum yield of total electron flow, which can be rewritten as:

$$\Phi_{e-} = \frac{4\left(\frac{F'q}{F'm} - b\right)}{k} \quad (5)$$

This relationship is assumed to hold in the presence and in the absence of photorespiration, where in the presence of photorespiration it can be used to calculate the total electron flow, which is the sum of electron flow to the reactions of carboxylation ( $J_C$ ) and oxygenation ( $J_O$ )<sup>35</sup>:

$$J_T = I * \Phi_{e-} = J_C + J_O \quad (6)$$

$J_C$  can be calculated by assuming 4 electrons are needed per  $\text{CO}_2$  for carboxylation:

$$J_C = 4(A + R_d + R_p) \quad (7)$$

where  $R_p$  is the amount of  $\text{CO}_2$  released by photorespiration.  $J_O$  can be calculated by assuming that 8 electrons are needed per  $\text{CO}_2$  released in photorespiration:

$$J_O = 8 * R_p \quad (8)$$

Combining the equations for  $J_T$ ,  $J_C$  and  $J_O$  then gives:

$$\begin{aligned} R_p &= \frac{J_T - 4(A + R_d)}{12} \\ J_C &= \frac{1}{3}[J_T + 8(A + R_d)] \\ J_O &= \frac{2}{3}[J_T - 4(A + R_d)] \end{aligned} \quad (9)$$

**Leaf transmission and reflectance.** Immediately following completion of the gas exchange measurements leaf absorbance ( $\alpha$ ) was measured from 400 to 700 nm, with a dual-channel spectrometer and integrating spheres (SpectroClip-JAZ-TR, Ocean Optics, Oxford, UK). Six spectral measurements of leaf transmittance and reflectance were made per leaf. The average transmittance ( $\tau$ ) and reflectance ( $R$ ) for each leaf was used to determine  $\alpha = (1 - R - \tau)$ . Total absorbance for the PAR spectrum was calculated. Total absorbed light was also calculated by combining the percentage of actinic light emitted by the blue (470 nm) and red (640 nm) LEDs in the leaf gas exchange chamber with  $\alpha$  for the peak wavelength of the two LED types<sup>36</sup>. The apparent maximum quantum yield ( $\Phi_{\max}$ ) was then recalculated to give the maximum absolute quantum yield, i.e. the net number of  $\text{CO}_2$  molecules absorbed by the leaf per photon absorbed:


$$\Phi_{\text{CO}_2, \max} = \frac{\Phi_{\max}}{\alpha} \quad (10)$$

**Intrinsic leaf water use efficiency.** Intrinsic leaf water use efficiency (LWUE) was calculated from gas exchange of  $\text{CO}_2$  and  $\text{H}_2\text{O}$  as the ratio of  $\text{CO}_2$  assimilation ( $A$ ) over stomatal conductance ( $g_s$ ) under 21%  $\text{O}_2$  at photon fluxes of 1000 and 2000  $\mu\text{mol m}^{-2} \text{s}^{-1}$ . Contrary to the water use efficiency calculated from  $A$  over transpiration ( $E$ )<sup>37</sup>,  $A/g_s$  is considered to be more realistic and comparable between studies, as it is not influenced by changes in leaf to air VPD in the leaf chamber<sup>38</sup>.

## Results

The light response of *A. donax* for  $A$  at high and low  $\text{O}_2$  showed an increase of  $A$  when  $\text{O}_2$  was lowered, compared to that at 21%  $\text{O}_2$ , indicating the reduction of  $A$  in air due to photorespiration (Fig. 1a). At both  $\text{O}_2$  concentrations,  $A$  did not reach a plateau, even at high photon flux. Both the light response of  $g_s$  and  $c_i$  were lowered when  $\text{O}_2$  concentration was reduced (Fig. 1b). The response of  $g_s$  to light largely scaled with  $A$ , while  $c_i$  reached a plateau above a photon flux of 600  $\mu\text{mol m}^{-2} \text{s}^{-1}$ . From the response of  $A$  to light, parameters were derived (Table 1). The  $\text{CO}_2$  assimilation rate at saturating photon flux ( $A_{\text{sat}}$ ) was remarkably high at both ambient (30.19  $\pm$  0.32  $\mu\text{mol m}^{-2} \text{s}^{-1}$ ) and low  $\text{O}_2$  concentration (34.86  $\pm$  1.2  $\mu\text{mol m}^{-2} \text{s}^{-1}$ ) whereas the predicted values of  $A$  at infinite photon flux density as described by the fitted hyperbola were 38.11  $\pm$  0.56  $\mu\text{mol m}^{-2} \text{s}^{-1}$  at ambient  $\text{O}_2$  and 43.45  $\pm$  2.08  $\mu\text{mol m}^{-2} \text{s}^{-1}$  at low  $\text{O}_2$  concentration. The maximum quantum yield of  $\text{CO}_2$  assimilation ( $\Phi_{\text{CO}_2, \max}$ ) at ambient and low  $\text{O}_2$  level was found to be 0.056 ( $\pm$  0.0029) and 0.072 ( $\pm$  0.0045) respectively, mol  $\text{CO}_2$  (mol absorbed photons)<sup>-1</sup>. The intrinsic leaf water use efficiency ( $A/g_s$ ) (LWUE) was 62.9 ( $\pm$  2.38)  $\mu\text{mol mol}^{-1}$  at sub-saturating photon flux (1000  $\mu\text{mol m}^{-2} \text{s}^{-1}$ ) and 66.0 ( $\pm$  1.97)  $\mu\text{mol mol}^{-1}$  at near-saturating photon flux (2000  $\mu\text{mol m}^{-2} \text{s}^{-1}$ ).

Under 21%  $\text{O}_2$ , the electron flow to oxygenation reactions of Rubisco ( $J_O$ ) and the rate of  $\text{CO}_2$  production by photorespiration ( $R_p$ ) increased with photon flux, until a plateau was reached at a photon flux of 1000  $\mu\text{mol m}^{-2} \text{s}^{-1}$



**Figure 1.** (a) Light response of assimilation rate for fully emerged *A. donax* leaves measured at 21% oxygen (closed symbols) and 2% oxygen (open symbols). (b)  $c_i$  (triangle) and  $g_s$  (square) at a  $c_a$  of  $400 \mu\text{mol mol}^{-1}$ , leaf temperature of  $25^\circ\text{C}$  and VPD of between 0.8 and 1.1 kPa (values are means and standard errors, for 5 different plants).


Photosynthetic light response curves	21% Oxygen	2% Oxygen
$A_{\text{sat}}$ ( $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ )	38.11 (0.56)	43.45 (2.08)
$\Phi_{\text{max}}$ (mol $\text{CO}_2$ mol photons $^{-1}$ )	0.050 (0.003)	0.064 (0.004)
$\Phi_{\text{CO}_2, \text{max}}$ (mol $\text{CO}_2$ mol photons $^{-1}$ )	0.056 (0.003)	0.072 (0.004)
$R_d$ ( $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ )	1.471 (0.15)	1.581 (0.16)
$\theta$ (—)	0.543 (0.06)	0.601 (0.08)
LCP ( $\mu\text{mol m}^{-2} \text{ s}^{-1}$ )	29.88 (2.14)	24.89 (1.63)

**Table 1.** Parameters derived from photosynthetic light response curves calculated with incident ( $I$ ) and absorbed photon flux ( $\alpha I$ ); replication and conditions of measurement as given in Fig. 1. mean, (SE),  $n = 5$ .


(Fig. 2.). However, the electron flow to carboxylation ( $J_C$ ) increased further with increasing photon flux and did not appear to reach a plateau (Fig. 2). It was estimated that *ca.* 25% of electron flow is accounted for by  $J_O$  under saturating light intensity. The ratio of  $\text{CO}_2$  production by photorespiration,  $R_p$ , to assimilation ( $A_{\text{sat}}$ ) was 0.216.

Total leaf absorbance from 400 nm to 700 nm was  $0.87 (+/- 4.0 \times 10^{-3}, n = 6)$ . Total leaf absorbance at the peak emitting wavelengths of red (640 nm) and blue (470 nm) was  $0.89$  (S.E.  $+/- 4.47 \times 10^{-3}, n = 6$ ) and used to calculate  $F_q'/F_m'$  against the quantum yield of  $\text{CO}_2$  assimilation ( $\Phi_{\text{CO}_2}$ ) on an absorbed light basis. The relation between  $F_q'/F_m'$  and  $\Phi_{\text{CO}_2}$  under high and low  $\text{O}_2$  showed significant positive correlations (Fig. 3). The difference






**Figure 2.** Electron flow to the reaction of carboxylation ( $J_c$ ) (open circle) and oxygenation ( $J_o$ ) (closed circle) and the rate of  $\text{CO}_2$  production by photorespiration ( $R_p$ ) (closed triangle) for fully emerged *A. donax* leaves. Values are corrected for leaf absorbance,  $0.89 (\pm 4.47 \times 10^{-3})$  and are calculated from light response of assimilation rate at a  $c_a$  of  $400 \mu\text{mol mol}^{-1}$ , leaf temperature of  $25^\circ\text{C}$  and VPD of between 0.8 and 1.1 kPa. (Values are means and standard errors, for 5 different plants).



**Figure 3.** Relationship between operating efficiency of PSII ( $F_q'/F_m'$ ) and quantum yield of  $\text{CO}_2$  assimilation ( $\Phi_{\text{CO}_2}$ ) for *A. donax* calculated from light response of assimilation rate, measured at 21% oxygen (closed circle) and 2% oxygen (open circle) and both at a  $c_a$  of  $400 \mu\text{mol mol}^{-1}$ , leaf temperature of  $25^\circ\text{C}$  and VPD of between 0.8 and 1.1 kPa. Values are corrected for leaf absorbance,  $0.89 (\pm 4.47 \times 10^{-3})$ , (values are means and standard errors, for 5 different plants).

in slope of the relation, representing the apparent electron demand to assimilate one  $\text{CO}_2$ , at ambient and low  $\text{O}_2$  concentration can be attributed to the presence or absence of photorespiration.

The response of  $A$  to  $c_i$  for *A. donax* leaves at saturating light intensity was determined at ambient  $\text{O}_2$  concentration (Fig. 4). Subsequently, parameters were derived to describe the  $A/c_i$  response and the stomatal limitation (Table 2). The maximum velocity of carboxylation by Rubisco ( $V_{c,\text{max}}$ ), was  $117.8 \mu\text{mol m}^{-2} \text{s}^{-1}$  and the maximum rate of electron transport ( $J_{\text{max}}$ ) was  $213.7 \mu\text{mol m}^{-2} \text{s}^{-1}$ . The  $\text{CO}_2$  compensation point,  $\Gamma$ , was  $44.2 \mu\text{mol mol}^{-1}$  and the limitation to  $A$  imposed by stomata and leaf boundary layer,  $l$ , was 0.25 (Table 2).



**Figure 4.** *A* vs *c<sub>i</sub>* response for *A. donax*. *A'*; the supply function of the estimated limitation placed on *A* by the stomata and leaf boundary layer; *A''* the supply function in the absence of any limitation to diffusion of CO<sub>2</sub> from the atmosphere to the site of carboxylation. Observations were measured at 21% oxygen on fully emerged leaves at a *I* of 1500 μmol m<sup>-2</sup> s<sup>-1</sup>, leaf temperature of 25 °C and VPD of 0.96 (0.01) kPa and *g<sub>s</sub>* 0.404 (0.016) mol m<sup>-2</sup> s<sup>-1</sup>. (Values are means and standard errors, for 5 different plants).

Photosynthetic intercellular-CO <sub>2</sub> response	21% Oxygen
<i>V<sub>c,max</sub></i> (μmol CO <sub>2</sub> m <sup>-2</sup> s <sup>-1</sup> )	117.8 (8.8)
<i>V<sub>c,max25</sub></i> (μmol CO <sub>2</sub> m <sup>-2</sup> s <sup>-1</sup> )	95.0 (6.9)
<i>J<sub>max</sub></i> (μmol CO <sub>2</sub> m <sup>-2</sup> s <sup>-1</sup> )	213.7 (21.2)
<i>J<sub>max25</sub></i> (μmol CO <sub>2</sub> m <sup>-2</sup> s <sup>-1</sup> )	184.0 (18.1)
Γ (μmol mol <sup>-1</sup> )	44.2 (0.95)
<i>l</i> (—)	0.25 (0.05)

**Table 2.** Parameters derived from photosynthetic intracellular CO<sub>2</sub> (*c<sub>i</sub>*) response curves; replication and conditions of measurement as given in Fig. 4. mean, (SE), *n* = 5.

Time	Leaf Surface	Stem	Middle	Tip	<i>F<sub>q</sub>'</i> / <i>F<sub>m</sub>'</i>
12:00	Abaxial	399 (61)	538 (72)	355 (25)	
12:00	Adaxial	293 (30)	422 (35)	326 (64)	0.63 (0.03)
16:00	Abaxial	231 (24)	373 (77)	341 (87)	
16:00	Adaxial	110 (10)	210 (18)	160 (28)	0.51 (0.08)

**Table 3.** Stomatal conductance (mmol m<sup>-2</sup> s<sup>-1</sup>) at mid and late afternoon, on abaxial and adaxial leaf surfaces with measurements along the leaf surface from the stem to the leaf tip. (Values are means and standard errors for 4 or 5 individual plants). The operating efficiency of PSII (*F<sub>q</sub>'*/*F<sub>m</sub>'*) at mid and late afternoon, on the middle adaxial leaf surfaces.

Soil water content between 20 cm and 40 cm soil depth averaged 0.14 m<sup>3</sup> m<sup>-3</sup> (+/− 0.01) across the sampling area over the sampling period, with an average soil temperature of 21 °C (+/− 0.9). Stomatal conductance of *A. donax*, measured on ad- and abaxial surfaces of leaves on three different positions along the leaf blade, were highest at noon (12:00) and declined in the afternoon (16:00, Table 3). The highest stomatal conductance was observed in the middle of the leaf at both time points and both surfaces. The greatest decline in stomatal conductance between noon and afternoon was observed in the middle of the leaf, on both adaxial and abaxial surfaces (Table 3). Even though there was a great decline in stomatal conductance, the operating efficiency of PSII (*F<sub>q</sub>'*/*F<sub>m</sub>'*) at noon and in the afternoon in the field did not decrease as much (Table 3). This implies that stomatal conductance was still not greatly limiting for photosynthesis during the afternoon.



## Discussion

Light-saturated capacity for photosynthetic  $\text{CO}_2$  uptake is determined by the maximum rate of carboxylation ( $V_{c,\max}$ ) and the maximum rate of electron transport, ( $J_{\max}$ ), depending on the inter-cellular  $\text{CO}_2$  concentration ( $c_i$ )<sup>20</sup>. Averaged across 109 C3 species,  $V_{c,\max}$  was  $64 \mu\text{mol m}^{-2} \text{s}^{-1}$  and  $J_{\max}$   $134 \mu\text{mol m}^{-2} \text{s}^{-1}$ <sup>39</sup>, this compares to values found here for *A. donax* of 117.8 and  $213.7 \mu\text{mol CO}_2 \text{m}^{-2} \text{s}^{-1}$ , respectively, which are almost double the C3 average. These values are for a natural and unfertilized stand. Typically in C3 crops these values will increase with nitrogen status<sup>40</sup>. However, the values found here for *A. donax* are high even compared to fertilized or N-fixing productive crops: *Phalaris arundinaceae* ( $83.9$  and  $150.6 \mu\text{mol m}^{-2} \text{s}^{-1}$ )<sup>41</sup>, *Triticum aestivum* ( $59$  and  $139 \mu\text{mol m}^{-2} \text{s}^{-1}$ )<sup>40</sup>, *Oryza sativa* ( $91$  and  $190 \mu\text{mol m}^{-2} \text{s}^{-1}$ )<sup>42</sup>, *Helianthus annuus* ( $29$  and  $Heinz\ walz \mu\text{mol m}^{-2} \text{s}^{-1}$ )<sup>43</sup>, *Glycine max* ( $83$  and  $160 \mu\text{mol m}^{-2} \text{s}^{-1}$ )<sup>44</sup>, and *Phaseolus vulgaris* ( $88$  and  $178 \mu\text{mol m}^{-2} \text{s}^{-1}$ )<sup>45</sup>. This highlights the high light-saturated maximum photosynthetic capacity of *A. donax*. Compared to other tall grass species considered as bioenergy sources, *A. donax* shows high  $A$  in full sunlight. Figure 1 shows an average  $A$  of  $30.2 \mu\text{mol m}^{-2} \text{s}^{-1}$  at  $25^\circ\text{C}$ , this compares to values at this temperature of  $28 \mu\text{mol m}^{-2} \text{s}^{-1}$  for *M. x giganteus*<sup>46</sup>,  $24 \mu\text{mol m}^{-2} \text{s}^{-1}$  in field grown stands of *Panicum virgatum* (switchgrass)<sup>16</sup>, and  $22 \mu\text{mol m}^{-2} \text{s}^{-1}$  for the C3 bioenergy grass, *Phalaris arundinacea* (Reed canary grass)<sup>47</sup>.

Is this  $A$  achieved by lower losses to photorespiration in *A. donax*? If we assume “average” Rubisco kinetic properties as described by Brooks and Farquhar<sup>48</sup> for spinach, then the  $\text{CO}_2$  photosynthetic compensation point ( $\Gamma^*$ ) would be  $42.5 \mu\text{mol mol}^{-1}$  at  $25^\circ\text{C}$ . Using this value, with the measured dark respiratory rate ( $R_d$ ) of  $2 \mu\text{mol m}^{-2} \text{s}^{-1}$ , average  $A$  of  $30.19 \mu\text{mol m}^{-2} \text{s}^{-1}$  and  $c_i$  of  $255 \mu\text{mol m}^{-2} \text{s}^{-1}$  at a photon flux of  $2000 \mu\text{mol m}^{-2} \text{s}^{-1}$  (Fig. 1), the photorespiratory rate of  $\text{CO}_2$  release ( $R_p$ ) would be  $6.3 \mu\text{mol m}^{-2} \text{s}^{-1}$ , following the equations of Farquhar *et al.*<sup>20</sup>. This calculated rate, based on properties of spinach, is very close to the rate derived here for *A. donax* by combining fluorescence and gas exchange measurements (Fig. 2) via the equations of Valentini *et al.*<sup>35</sup>. This suggests that as a proportion of net photosynthesis, photorespiratory loss of carbon is just as great in *A. donax* as in other C3 species.

Is the rate of  $\text{CO}_2$  assimilation in limiting light high in *A. donax*? Light limited photosynthesis is determined by the efficiency with which the leaf can absorb incident light ( $\alpha$ ), and the maximum absolute quantum yield of  $\text{CO}_2$  assimilation ( $\Phi_{\text{CO}_2,\max}$ ), i.e. the initial slope of the response of  $A$  to  $I$ , corrected for absorbance ( $\alpha$ ). The value is therefore the maximum ratio of net absorbed  $\text{CO}_2$  molecules to absorbed photons. The  $\Phi_{\text{CO}_2,\max}$  determined for *A. donax* here in normal air at  $25^\circ\text{C}$  was  $0.056 \text{ mol mol}^{-1}$ . Osborne and Garrett<sup>49</sup> similarly measured  $\Phi_{\text{CO}_2,\max}$  by combining gas exchange and integrating sphere measurements across a range of C3 herbage grasses and cereals, covering different ploidy levels and cultivars in normal air and at  $25^\circ\text{C}$ . They reported an average  $\Phi_{\text{CO}_2,\max}$  of  $0.051$ , with a range of  $0.047$  to  $0.055$  across these C3 grasses. So this does place the values here for *A. donax* at the upper end of this range and  $8\%$  higher than the average. The realized efficiency of light limited photosynthesis will be the product of  $\Phi_{\text{CO}_2,\max}$  and the absorbance of the leaf ( $\alpha$ ). In surveying a wide range of healthy leaves of C3 species from tropical to polar habitats, Long *et al.*<sup>50</sup>, found an average  $\alpha$  of  $0.80$ , and a range across species from  $0.65$  to  $0.90$ , similarly measured in an integrating sphere. By comparison, the  $\alpha$  for *A. donax* of  $0.89$  therefore appears at the upper end of this range and  $14\%$  higher above the average. The product of the indicated above average  $\Phi_{\text{CO}_2,\max}$  and  $\alpha$  therefore represents a  $24\%$  increase in  $\text{CO}_2$  uptake per unit incident light, under light-limiting conditions.

Does *A. donax* achieve high photosynthetic rates by minimizing stomatal limitation? The stomatal limitation imposed on photosynthesis,  $l$ , for *A. donax* found in this study ( $0.25$  or  $25\%$ ) was higher than for other C3 plants, which varied from  $0.137$  to  $0.217$ <sup>51–54</sup>. This indicates that *A. donax* does not achieve its high photosynthetic rates through a high stomatal conductance. Indeed  $c_i/c_a$  at full sunlight ( $2000 \mu\text{mol m}^{-2} \text{s}^{-1}$ ) as a measure of the balance between stomatal conductance and assimilation was  $0.64$ . This is almost  $11\%$  less than the average  $c_i/c_a$  of  $0.72$  for a range of C3 species<sup>19</sup>. A lower  $c_i/c_a$  requires a lower stomatal conductance relative to the photosynthetic rate, indicating a higher leaf level water use efficiency for a given leaf-air water vapour pressure deficit than the average for C3 species. So, while stomatal conductance may appear high (Fig. 1b), the low values of  $l$  and  $c_i/c_a$  in fact show that conductance is low relative to the high rates of leaf  $\text{CO}_2$  uptakes for a C3 plant. Nevertheless its high light-saturated and light-limited photosynthetic rates can still only be achieved at the expense of considerable transpiration. The relatively high stomatal conductance maintained into the late afternoon (Table 3) suggests that the plant may be able to tap deep water resources even in its native Mediterranean in the late summer. The LWUE of *A. donax* in this study ( $62.9$  to  $66.0 \mu\text{mol mol}^{-1}$ ) was higher than generally found for herbaceous species ( $43 \mu\text{mol mol}^{-1}$ ) and more similar to LWUE found for evergreen shrubs and deciduous trees ( $64$  and  $66 \mu\text{mol mol}^{-1}$ , respectively)<sup>55</sup>. However, the LWUE of *A. donax* was still much lower than for C4 species such as *Miscanthus* and switchgrass ( $115$  and  $107 \mu\text{mol mol}^{-1}$ )<sup>16</sup>. The high productivity of *A. donax* does resemble and exceeds that of C4 plants, however its LWUE is much lower and clearly C3-like. This supports the notion that, as discussed above, *A. donax* is able to achieve its high photosynthetic rates with substantial transpiration, but is still more efficient than most C3 species. This is interesting when considering the diversity of habitat distribution that *A. donax* is found in, which ranges from very wet loam to relatively dry sandy soils.

Overall this study has found that the photosynthetic capacity of *A. donax* in full sunlight is high compared to other C3 species, and comparable to C4 bioenergy grasses. This is not the result of lower photorespiratory rates, but rather a high capacity for both RuBP-limited and RuBP-saturated photosynthesis, as evidenced by values of both  $V_{c,\max}$  and  $J_{\max}$  that are near double the average for C3 species. High photosynthetic rates were not achieved through a high stomatal conductance, in fact stomatal limitation was found to be greater not less than in other C3 species. Light-limited photosynthesis, which will determine carbon uptake during periods of low light flux, around dawn and dusk, and during heavy cloud, as well as in the lower canopy may be aided by relatively high maximum quantum yields of  $\text{CO}_2$  assimilation and high leaf absorbances.

## References

1. EPA. *Supplemental Final Rule for New Qualifying Renewable Fuels under the RFS Program*. Environ. Prot. Agency - Off. Transp. Air Qual. (2013).
2. Lewandowski, I., Scurlock, J. M. O., Lindvall, E. & Christou, M. The development and current status of perennial rhizomatous grasses as energy crops in the US and Europe. *Biomass and Bioenergy* **25**, 335–361 (2003).
3. Pilu, R. Giant reed (*Arundo donax* L.): A weed plant or a promising energy crop? *African J. Biotechnol.* **11**, 9163–9174 (2012).
4. Di Nasso, N., Roncucci, N. & Bonari, E. Seasonal Dynamics of Aboveground and Belowground Biomass and Nutrient Accumulation and Remobilization in Giant Reed (*Arundo donax* L.): A Three-Year Study on Marginal Land. *BioEnergy Res.* **6**, 725–736 (2013).
5. Mariani, C. *et al.* Origin, diffusion and reproduction of the giant reed (*Arundo donax* L.): a promising weedy energy crop. *Ann. Appl. Biol.* **157**, 191–202 (2010).
6. Angelini, L. G., Ceccarini, L., Nasso, N. & Bonari, E. Comparison of *Arundo donax* L. and *Miscanthus x giganteus* in a long-term field experiment in Central Italy: Analysis of productive characteristics and energy balance. *Biomass and Bioenergy* **33**, 635–643 (2009).
7. Heaton, E. A quantitative review comparing the yields of two candidate C<sub>4</sub> perennial biomass crops in relation to nitrogen, temperature and water. *Biomass and Bioenergy* **27**, 21–30 (2004).
8. Borin, M. *et al.* Biomass production and N balance of giant reed (*Arundo donax* L.) under high water and N input in Mediterranean environments. *Eur. J. Agron.* **51**, 117–119 (2013).
9. Ping, H., Bransby, D. I. & van Santen, E. Long-term biomass yields of giant reed, mimosa and switchgrass in Alabama. *Biofuels, Bioprod. Biorefining* **8**, 59–66 (2014).
10. Ceotto, E. *et al.* Comparing solar radiation interception and use efficiency for the energy crops giant reed (*Arundo donax* L.) and sweet sorghum (*Sorghum bicolor* L. Moench). *F. Crop. Res.* **149**, 159–166 (2013).
11. Drewry, D. T., Kumar, P. & Long, S. P. Simultaneous improvement in productivity, water use, and albedo through crop structural modification. *Glob. Chang. Biol.* **20**, 1955–67 (2014).
12. Arundale, R. *et al.* Yields of *Miscanthus x giganteus* and *Panicum virgatum* decline with stand age in the Midwestern USA. *Glob. Chang. Biol. - Bioenergy* **6**, 1–13 (2014).
13. Mann, J. J., Kyser, G. B., Barney, J. N. & DiTomaso, J. M. Assessment of Aboveground and Belowground Vegetative Fragments as Propagules in the Bioenergy Crops *Arundo donax* and *Miscanthus x giganteus*. *bioenerg* **6**, 688–698 (2013).
14. Papazoglou, E. G., Karantounias, G. A., Vemmos, S. N. & Bouranis, D. L. Photosynthesis and growth responses of giant reed (*Arundo donax* L.) to the heavy metals Cd and Ni. In *Environ. Int.* **31**, 243–249 (2005).
15. Rossa, B., Tuffers, A. V., Naidoo, G. & Willert, D. J. Von. *Arundo donax* L. (Poaceae) - a C<sub>3</sub> Species with Unusually High Photosynthetic Capacity. *Bot. Acta* **111**, 216–221 (1998).
16. Dohleman, F. G., Heaton, E. A., Leakey, A. D. B. & Long, S. P. Does greater leaf-level photosynthesis explain the larger solar energy conversion efficiency of *Miscanthus* relative to switchgrass? *Plant. Cell Environ.* **32**, 1525–37 (2009).
17. Dohleman, F. G. & Long, S. P. More productive than maize in the Midwest. How does *Miscanthus* do it? *Plant Physiol. Preview*, 109.139162 (2009).
18. Long, S. P. & Spence, A. K. Toward cool C<sub>4</sub> crops. *Annu. Rev. Plant Biol.* **64**, 701–22 (2013).
19. Long, S. P., Ainsworth, E. A., Rogers, A. & Ort, D. R. Rising Atmospheric Carbon Dioxide: Plants FACE the Future. *Annu. Rev. Plant Biol.* **55**, 591–628 (2004).
20. Farquhar, G. D., Caemmerer, S., Von & Berry, J. A. A Biochemical Model of Photosynthetic CO<sub>2</sub> Assimilation in Leaves of C<sub>3</sub> Species. *Planta* **149**, 78–90 (1980).
21. Wang, D. *et al.* A physiological and biophysical model of coppice willow (*Salix* spp.) production yields for the contiguous USA in current and future climate scenarios. *Plant. Cell Environ.* **38**, 1850–1865 (2015).
22. ISRIC. (2014). at <www.isric.org>.
23. Monteith, J. L., Campbell, G. S. & Potter, E. A. Theory and performance of a dynamic diffusion porometer. *Agric. For. Meteorol.* **44**, 27–38 (1988).
24. Potter, L., Bingham, M. G., Baker, M. G. & Long, S. P. The potential of two perennial C<sub>4</sub> grasses and a perennial C<sub>4</sub> sedge as ligno-cellulosic fuel crops in N. W. Europe crop establishment and yields in E. England. *Ann. Bot.* **76**, 513–520 (1995).
25. Baker, N. R. Chlorophyll Fluorescence: A Probe of Photosynthesis *In Vivo*. *Annu. Rev. Plant Biol.* **59**, 89–113 (2008).
26. Murchie, E. H. & Lawson, T. Chlorophyll fluorescence analysis: a guide to good practice and understanding some new applications. *J. Exp. Bot.* **64**, 3983–98 (2013).
27. Bernacchi, C. J., Singsaas, E. L., Pimentel, C., Portis Jr., A. R. & Long, S. P. Improved temperature response functions for models of Rubisco-limited photosynthesis. *Plant, Cell Environ.* **24**, 253–259 (2001).
28. Bernacchi, C. J., Pimentel, C. & Long, S. P. *In vivo* temperature response functions of parameters required to model RuBP-limited photosynthesis. *Plant. Cell Environ.* **26**, 1419–1430 (2003).
29. Long, S. P. & Bernacchi, C. J. Gas exchange measurements, what can they tell us about the underlying limitations to photosynthesis? Procedures and sources of error. *J. Exp. Bot.* **54**, 2393–2401 (2003).
30. Genty, B., Briantais, J.-M. & Baker, N. R. The relationship between the quantum yield of photosynthetic electron transport and quenching of chlorophyll fluorescence. *Biochim. Biophys. Acta* **990**, 87–92 (1989).
31. Marshall, B. & Biscoe, P. V. A Model for C<sub>3</sub> Leaves Describing the Dependence of Net Photosynthesis on Irradiance. *J. Exp. Bot.* **31**, 29–39 (1980).
32. Genty, B., Harbinson, J. & Baker, N. Relative quantum efficiencies of the two photosystems of leaves in photorespiratory and non-respiratory conditions. *Plant Physiol. Biochem.* **28**, 1–10 (1990).
33. Edwards, G. E. & Baker, N. R. Can CO<sub>2</sub> assimilation in maize leaves be predicted accurately from chlorophyll fluorescence analysis? *Photosynth. Res.* **37**, 89–102 (1993).
34. Lawson, T. *et al.* C<sub>3</sub> photosynthesis in the desert plant *Rhazya stricta* is fully functional at high temperatures and light intensities. *New Phytol.* **201**, 862–873 (2014).
35. Valentini, R., Epron, D., Angelis, P. D. E., Matteucci, G. & Dreyer, E. *In situ* estimation of net CO<sub>2</sub> assimilation, photosynthetic electron flow and photorespiration in Turkey oak (*Q. cerris* L.) leaves: diurnal cycles under different levels of water supply. *Plant Cell Environ.* **18**, 631–640 (1995).
36. Naidu, S. L. & Long, S. P. Potential mechanisms of low temperature tolerance of C<sub>4</sub> photosynthesis in *Miscanthus x giganteus*: an *in vivo* analysis. *Planta* **220**, 145–155 (2004).
37. Penman, H. L. & Schofield, R. K. Some physical aspects of assimilation and transpiration. in *Symp. Soc. Exp. Biol* **5**, 115–129 (1951).
38. Bacon, M. *Water use efficiency in plant biology*. (John Wiley & Sons, 2009).
39. Wullschlegel, S. D. Biochemical Limitations to Carbon Assimilation in C<sub>3</sub> Plants—A Retrospective Analysis of the A/C<sub>i</sub> Curves from 109 Species. *J. Exp. Bot.* **44**, 907–920 (1993).
40. Evans, J. R. & Farquhar, G. D. In *Model. Crop Photosynth. Biochem. to canopy* (Boote, K. & Loomis, R.) 1–15 (American Society of Agronomy, Inc, 1991).
41. He, Z., Bentley, L. P. & Holaday, A. S. Greater seasonal carbon gain across a broad temperature range contributes to the invasive potential of *Phalaris arundinacea* (Poaceae; reed canary grass) over the native sedge *Carex stricta* (Cyperaceae). *Am. J. Bot.* **98**, 20–30 (2011).

42. Ziska, L. H. & Teramura, A. H. CO<sub>2</sub> Enhancement of Growth and Photosynthesis in Rice (*Oryza sativa*) : Modification by Increased Ultraviolet-B Radiation. *Plant Physiol.* **99**, 473–481 (1992).
43. Siebke, K. *et al.* Control of photosynthesis in leaves as revealed by rapid gas exchange and measurements of the assimilatory force FA. *Planta* **182**, 513–522 (1990).
44. Harley, P. C. & Sharkey, T. D. An improved model of C<sub>3</sub> photosynthesis at high CO<sub>2</sub>: Reversed O<sub>2</sub> sensitivity explained by lack of glycerate reentry into the chloroplast. *Photosynth. Res.* **27**, 169–178 (1991).
45. Sage, R. F., Sharkey, T. D. & Seemann, J. R. Acclimation of Photosynthesis to Elevated CO<sub>2</sub> in Five C<sub>3</sub> Species. *Plant Physiol.* **89**, 590–596 (1989).
46. Naidu, S. L., Moose, S. P., Al-Shoaibi, A. K., Raines, C. A. & Long, S. P. Cold Tolerance of C<sub>4</sub> photosynthesis in *Miscanthus x giganteus*: *Adaptation in Amounts and Sequence of C<sub>4</sub> Photosynthetic Enzymes*. **132**, 1688–1697 (2003).
47. Ge, Z. M. *et al.* Acclimation of photosynthesis in a boreal grass (*Phalaris arundinacea* L.) under different temperature, CO<sub>2</sub>, and soil water regimes. *Photosynthetica* **50**, 141–151 (2012).
48. Brooks, A. & Farquhar, G. D. Effect of temperature on the CO<sub>2</sub>/O<sub>2</sub> specificity of ribulose-1,5-bisphosphate carboxylase/oxygenase and the rate of respiration in the light. *Planta* **165**, 397–406 (1985).
49. Osborne, B. A. & Garrett, M. K. Quantum yields for CO<sub>2</sub> uptake in some diploid and tetraploid plant species. *Plant. Cell Environ.* **6**, 135–144 (1983).
50. Long, S. P., Postl, W. F. & Bolh r-Nordenkamp, H. R. Quantum yields for uptake of carbon-dioxide in C-3 vascular plants of contrasting habitats and taxonomic groupings. *Planta* **189**, 226–234 (1993).
51. Maroco, J. P., Rodrigues, M. L., Lopes, C. & Chaves, M. M. Limitations to leaf photosynthesis in field-grown grapevine under drought - metabolic and modelling approaches. *Funct. Plant Biol.* **29**, 451 (2002).
52. Bernacchi, C. J., Morgan, P. B., Ort, D. R. & Long, S. P. The growth of soybean under free air [CO<sub>2</sub>] enrichment (FACE) stimulates photosynthesis while decreasing *in vivo* Rubisco capacity. *Planta* **220**, 434–446 (2005).
53. Ripley, B. S., Gilbert, M. E., Ibrahim, D. G. & Osborne, C. P. Drought constraints on C<sub>4</sub> photosynthesis: Stomatal and metabolic limitations in C<sub>3</sub> and C<sub>4</sub> subspecies of *Alloteropsis semialata*. *J. Exp. Bot.* **58**, 1351–1363 (2007).
54. Hu, L., Wang, Z. & Huang, B. Diffusion limitations and metabolic factors associated with inhibition and recovery of photosynthesis from drought stress in a C<sub>3</sub> perennial grass species. *Physiol. Plant.* **139**, 93–106 (2010).
55. Medrano, H., Flexas, J. & Galm s, J. Variability in water use efficiency at the leaf level among Mediterranean plants with different growth forms. *Plant Soil* **317**, 17–29 (2009).

## Acknowledgements

The authors thank the Society of Experimental Biology Plant Environmental Physiology Group (PEPg) Techniques Workshop, Lisbon 2014 for bringing the authors together. This was also made possibly by financial support from NERC Advanced training award; NE/M006395/1, the British Ecology Society and the Journal of Experimental Botany. We also thank ADC BioScientific Ltd, Decagon Devices, Delta-T Devices Ltd, Dynamax Inc., LI-COR Biosciences UK Ltd, Ocean Optics and Heinz Walz GmbH for providing equipment, and site manager, Benjamin Richter, of Quinta de S o Pedro Study Centre, Lisbon, Portugal, for logistical support.

## Author Contributions

R.W. and S.D. conceived and designed the study. R.W., S.D., S.L., J.K., J.M., A.L., K.S., T.D., T.L., T.P. and S.B. all contributed to data collection and analysis. R.W. and S.D. produced the final analysis. R.W., S.D. and S.L. primarily led manuscript preparation. All authors reviewed the final manuscript.

## Additional Information

**Supplementary information** accompanies this paper at <http://www.nature.com/srep>

**Competing financial interests:** The authors declare no competing financial interests.

**How to cite this article:** Webster, R. J. *et al.* High C<sub>3</sub> photosynthetic capacity and high intrinsic water use efficiency underlies the high productivity of the bioenergy grass *Arundo donax*. *Sci. Rep.* **6**, 20694; doi: 10.1038/srep20694 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>